

# Progress of the 1/12° Global HYCOM Effort

E. Joseph Metzger and Alan J. Wallcraft

HYCOM Consortium Meeting

6-8 December 2005

RSMAS, Miami FL

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>DEC 2005</b>		2. REPORT TYPE		3. DATES COVERED <b>00-00-2005 to 00-00-2005</b>	
4. TITLE AND SUBTITLE <b>Progress of the 1/12degree Global HYCOM Effort</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Naval Research Laboratory,Stennis Space Center,MS,39529</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES <b>20</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

# 1/12° Global HYCOM: Initial Development and Evaluation

E.J. Metzger, A.J. Wallcraft, H.E. Hurlburt,  
E.P. Chassignet and W.J. Schmitz, Jr.

Ocean Sciences Meeting  
20-24 February 2006  
Honolulu, HI


# HYCOM Long-term Goals for Operational Ocean Prediction

- 1/12° fully global ocean prediction system transitioned to NAVO in 2007
  - Include shallow water, minimum depth 5 m
  - Coupled sea-ice model (LANL CICE)
- Increase to 1/25° resolution globally by the end of the decade
  - Optimal resolution for basin-scale
  - Boundary conditions for coastal models

# Global HYCOM Configuration

- Horizontal grid: 1/12° equatorial resolution
  - 4500 x 3298 grid points, 6.5 km spacing on average, 3.5 km at pole
- Mercator 79°S to 47°N, then Arctic dipole patch
- Vertical coordinate surfaces: 26-28 for  $\sigma_0$ , 32 for  $\sigma_2^*$
- KPP and GISS mixed layer models
- Thermodynamic (energy loan) sea-ice model
- Surface forcing: wind stress, wind speed, thermal forcing, precipitation, relaxation to climatological SSS
- Monthly river runoff (986 rivers)
- Initialize from January climatology (GDEM3) T and S, then SSS relaxation from PHC 3.0
  - No subsurface relaxation to climatology

# 1/12° Global HYCOM snapshot: SSH and ice (gray)



- Running at NAVO under DoD Challenge
- 190K CPU hrs/model year on 784 CPUs
- 7.2 TB/model year for daily 3-D output


# 1/12° Global HYCOM Experiments

- ECMWF Reanalysis (ERA15) climatological wind and thermal forcing
  - Annual bias corrections to air temperature (ERA40), radiative fluxes (ISCCP) and precipitation (GPCP)
- $\sigma_0$  simulations:
  - 26-layers with KPP for 6 model years
  - 28-layers with GISS for 3 model years
- $\sigma_2^*$  simulations:
  - 32-layers with GISS for 9 model years



# Initial 1/12° Global HYCOM $\sigma_0$ Simulation



- Major shortcomings:
  - Poor simulation of both Gulf Stream and Kuroshio
  - Poor representation of tropical current systems
  - Unrealistic transport at key locations:
    - Florida Straits (23 Sv vs. ~32 Sv) [simulated vs. observed]
    - Drake Passage (91 Sv vs. ~134 Sv)
    - Pacific to Indian Ocean Throughflow (22 Sv vs. ~10 Sv)





# Improved 1/12° Global HYCOM $\sigma_\theta$ Simulation

- Modifications
  - Added two layers (26  $\rightarrow$  28) and changed layer structure
  - Increased eddy viscosity:  $A = 30 \text{ m}^2/\text{s}$  constant everywhere
  - Increased Smagorinsky diffusion:  $.05 \rightarrow .1$
  - KPP  $\rightarrow$  GISS



# Improvements in Tropical Pacific Current Structure


Observations

Original 26-layer  $\sigma_0$


Modified 28-layer  $\sigma_0$

Zonal velocity along the equator


Zonal velocity at 140°W



# 1/12° Global HYCOM Mean Gulf Stream And Kuroshio Pathways



Latest  $\sigma_2^*$  simulation




Mean over four model years

ERA15 climatological wind & thermal forcing




## The Atlantic subpolar gyre generally looks good




11

# Global SSH Variability



Oct 92 – Nov 98 SSH variability based on T/P, ERS-1 and ERS-2 altimeters (from Collecte, Localisation, Satellites (CLS))




SSH variability from 1/12° global HYCOM  $\sigma_2^*$  with climatological wind and thermal forcing





# Mediterranean Sea Outflow


## Salinity section at 36°N






# Atlantic Meridional Overturning Circulation

$\sigma_0$  vs.  $\sigma_2^*$




# Atlantic Meridional Overturning Circulation




# Velocity Cross-section Along Luzon Strait

Sb-ADCP data (top) vs. 1/12° global HYCOM (bottom) in the upper 300 m  
Section along 21°N between 118.5°E and 124.0°E



Cross-section overlaid on mean currents and speed



Sb-ADCP data from Liang et al. (2003, DSR Pt. II)  
Mean from HYCOM with ERA15 wind and thermal forcing  
No ocean data assimilation in HYCOM





# Vertical Structure in the Equatorial Pacific


Observations

32-layer  $\sigma_2^*$

Modified 28-layer  $\sigma_0$





# Deep Flow Through Samoan Passage



Observed mean northward  
transport below 4000 m = 6.0 Sv  
Rudnick (1997, JGR)

HYCOM  $\sigma_2^*$  in layers 27-32 = 9.3 Sv



# Transport Comparisons at Key Locations

Section	Obs.	Orig. $\sigma_0$	Mod. $\sigma_0$	Orig. $\sigma_2^*$	Mod. $\sigma_2^*$
PCM-1	23	24.8	24.7	25.5	26.4
Bering Strait	1	1.1	1.1	1.1	1.1
PACIO TF	-10	-21.8	-24.1	-18.2	-17.0
STACS	30-34	23.2	23.1	22.9	24.0
Yucatan Channel	23-27	23.3	21.6	21.8	22.0
Denmark Strait	-2.9	-2.9	-2.3	-2.3	-2.9
Drake Passage	134	91.3	96.2	152.4	146.4

Modified  $\sigma_2^*$  experiment uses a new topography with sill depth refinements in the IAS, Indonesian Seas, etc.



# Future Work (FY06)

- Ten May 2001 – June 2002 assimilative runs in FY06
  - Time period with three satellite altimeters
  - Five with bi-weekly 30-day forecasts
- Near real-time nowcast/forecast starting in mid-FY06
- Interannual non-assimilative case:
  - 1995-present using NOGAPS
- Coupling with LANL CICE via ESMF